

Review Article

Anredera cordifolia (Binahong) in Hypertension Management: Phytochemical Basis, Cardiovascular Mechanisms, and Current Evidence

La Ode Alifariki^{1,*}, Dewi Nopiska Lilis²

¹Department of Nursing, Medical Faculty, Halu Oleo University, Kendari, Indonesia

²Department of Midwifery, Health Polytechnic of the Ministry of Health Jambi, Jambi, Indonesia

Article information

Received: June 05, 2026

Accepted: July 22, 2026

Published: July 31, 2026

*Corresponding author:

La Ode Alifariki,

E-mail: ners_riki@yahoo.co.id

Keywords: *Anredera cordifolia*; Binahong; Hypertension; Herbal Medicine; Blood pressure; Endothelial function; Phytochemicals

Cite this article: La Ode Alifariki, and Dewi Nopiska Lilis. *Anredera cordifolia* (Binahong) in Hypertension Management: Phytochemical Basis, Cardiovascular Mechanisms, and Current Evidence. J Biosci Public Health. 2026;2(4):293-312. doi:10.5455/JBPH.2026.14

Copyright © The Author(s):

This is an open access article under the (CC BY 4.0) license(<https://creativecommons.org/licenses/by/4.0/deed.en>).

ABSTRACT

Background: Hypertension remains one of the leading global public health challenges and a major contributor to cardiovascular morbidity and mortality. Despite the availability of effective pharmacological therapies, blood pressure control remains suboptimal in many populations, leading to growing interest in complementary and herbal approaches. *Anredera cordifolia* (Binahong) is traditionally used for various health conditions and has attracted attention for its potential cardiovascular benefits. **Objective:** Aimed to synthesize and critically evaluate current evidence regarding the potential role of Binahong leaf decoction in hypertension management, with emphasis on its phytochemical characteristics, biological mechanisms, safety profile, and implications for clinical practice. **Methods:** A narrative review was conducted using literature retrieved from PubMed, Scopus, Web of Science, ScienceDirect, and Google Scholar. A total of 11 studies were included in the final synthesis, comprising phytochemical investigations, antioxidant studies, mechanistic pharmacological studies, animal experiments, and comparative clinical studies involving herbal antihypertensive interventions. **Results:** Evidence indicates that *A. cordifolia* contains bioactive compounds, including flavonoids, polyphenols, saponins, alkaloids, and terpenoids, with antioxidant and vasoprotective properties. Experimental studies suggest potential antihypertensive effects through antioxidant activity, endothelial protection, vasodilation, and moderate angiotensin-converting enzyme inhibition. However, direct clinical evidence remains scarce, with available human studies limited by small sample sizes, methodological heterogeneity, and non-randomized designs. Current data are therefore insufficient to establish clinical efficacy or standardized therapeutic protocols. **Conclusions:** *Anredera cordifolia* demonstrates mechanisms that may support blood pressure regulation and cardiovascular protection. Well-designed experimental studies and randomized controlled trials are required to determine its effectiveness, safety, optimal dosage, and potential role in integrative hypertension care.

1. INTRODUCTION

Hypertension remains one of the most significant global public health challenges and is a leading modifiable risk factor for cardiovascular morbidity and mortality [1, 2]. Persistent elevation of blood pressure contributes substantially to the development of coronary artery disease, stroke, heart failure, chronic kidney disease, peripheral vascular disease, and premature death [3]. Despite considerable advances in prevention, diagnosis, and treatment, hypertension continues to affect more than one billion individuals worldwide and is responsible for millions of deaths annually. The burden of hypertension is particularly pronounced in low- and middle-income countries, where healthcare systems frequently face challenges related to screening, diagnosis, treatment accessibility, medication adherence, and long-term disease management [4, 5].

The increasing prevalence of hypertension has been attributed to population aging, urbanization, sedentary lifestyles, unhealthy dietary patterns, obesity, tobacco use, excessive alcohol consumption, and psychosocial stress [6]. Although pharmacological therapies such as angiotensin-converting enzyme (ACE) inhibitors, angiotensin receptor blockers, calcium channel blockers, beta-blockers, and diuretics have demonstrated efficacy in reducing cardiovascular risk, achieving optimal blood pressure control remains a persistent challenge. Studies have shown that a substantial proportion of hypertensive patients fail to achieve target blood pressure levels due to poor medication adherence, adverse drug reactions, treatment fatigue, economic barriers, and limited access to healthcare services. Consequently, interest in complementary and alternative therapeutic approaches has increased substantially over the past decade [7].

Herbal medicine constitutes one of the most widely used forms of complementary therapy worldwide [3]. Cultural acceptance, perceived safety, affordability, accessibility, and longstanding traditional use drive the growing utilization of medicinal plants. In many Asian countries, herbal preparations continue to play an important role in primary healthcare and chronic disease management. Furthermore, increasing scientific attention has been directed toward identifying plant-derived bioactive compounds with potential antihypertensive, antioxidant, anti-inflammatory, and cardioprotective properties. Such investigations are particularly relevant because hypertension is increasingly recognized as a multifactorial disorder involving oxidative stress, endothelial dysfunction, chronic inflammation, vascular remodeling, sympathetic overactivity, and dysregulation of the renin–angiotensin–aldosterone system (RAAS) [8].

Among the medicinal plants commonly utilized in Southeast Asia, *Anredera cordifolia* (Ten.) Steenis, locally known as Binahong, has gained considerable attention due to its diverse pharmacological properties. Binahong is a perennial climbing plant belonging to the Basellaceae family and is widely distributed throughout tropical and subtropical regions. Although native to South America, the plant has become extensively cultivated in Indonesia

and several Asian countries, where it is traditionally used for wound healing, gastrointestinal disorders, diabetes management, inflammatory conditions, and cardiovascular health promotion. The leaves represent the most frequently utilized part of the plant and are commonly prepared as decoctions, infusions, extracts, or topical formulations [9, 10].

Traditionally, Binahong leaves are prepared by boiling about 5-10 g of fresh leaves in 200-300 mL of water for 10-15 minutes before oral consumption. However, preparation methods differ considerably among communities with respect to the amount of plant material, water volume, boiling duration, and frequency of administration. To date, no standardized decoction protocol has been established. This variability may substantially influence the extraction efficiency and concentration of bioactive compounds, thereby affecting the pharmacological activity and reproducibility of research findings. Consequently, differences in preparation methods should be considered when interpreting the available evidence regarding the antihypertensive potential of Binahong [11, 12].

Phytochemical investigations have revealed that Binahong leaves contain a wide range of bioactive constituents, including flavonoids, polyphenols, saponins, alkaloids, terpenoids, tannins, glycosides, and various antioxidant compounds [12-14]. These secondary metabolites have attracted scientific interest because of their potential roles in cardiovascular protection. Flavonoids and polyphenols, in particular, are known to exhibit potent antioxidant activities capable of reducing reactive oxygen species, enhancing nitric oxide bioavailability, and protecting endothelial function. Endothelial dysfunction is recognized as a central mechanism in hypertension pathogenesis, and interventions that improve endothelial health may contribute to blood pressure reduction and cardiovascular risk mitigation [15].

Emerging evidence suggests that Binahong may exert antihypertensive effects through multiple complementary pathways. Experimental studies have demonstrated vasodilatory activity in isolated vascular tissues, indicating a direct influence on vascular tone regulation. In addition, several investigations have reported moderate ACE inhibitory activity, suggesting a potential role in modulating the RAAS pathway, which is critically involved in blood pressure regulation [16]. Other studies have highlighted the antioxidant and anti-inflammatory properties of Binahong, which may attenuate oxidative stress, reduce vascular inflammation, and improve arterial function. Collectively, these findings provide biological plausibility for the traditional use of Binahong in hypertension management [10]. However, clinical investigations are mostly small-scale, quasi-experimental studies with methodological limitations, including small sample sizes, absence of control groups, short intervention durations, and lack of standardized preparation methods [17]. Few studies have directly evaluated blood pressure outcomes, and even fewer have investigated the effects of traditionally prepared Binahong leaf decoctions in human populations. Another important limitation is the absence of large-scale randomized controlled trials examining the efficacy and safety of Binahong in hypertensive populations. Questions regarding optimal dosage, preparation

methods, treatment duration, long-term safety, potential toxicity, and herb-drug interactions remain largely unanswered.

Therefore, this narrative review aims to synthesize and critically evaluate the current evidence regarding the potential role of Binahong (*Anredera cordifolia*) leaf decoction in hypertension management. Specifically, this review examines the phytochemical composition of Binahong, proposed antihypertensive mechanisms, evidence from experimental and human studies, preparation methods, safety considerations, and implications for clinical practice and future research. By integrating available evidence, this review seeks to provide a comprehensive understanding of the therapeutic potential and current limitations of Binahong as a complementary approach to hypertension management.

2. METHODOLOGY

2.1. Review Design

This study was conducted as a narrative review. Although a structured literature search and predefined eligibility criteria were applied to improve transparency and comprehensiveness, the review did not follow the full methodological requirements of a systematic review, such as protocol registration, formal risk-of-bias assessment, or quantitative evidence synthesis. This study employed a narrative review design to comprehensively synthesize and critically evaluate the available evidence regarding the use of Binahong (*Anredera cordifolia*) leaf preparations, particularly leaf decoctions, for hypertension management [12]. A narrative review methodology was selected because the available literature is relatively limited and heterogeneous, consisting of phytochemical investigations, mechanistic pharmacology studies, experimental animal research, ex vivo vascular studies, and a small number of human intervention studies. The diversity of study designs and outcome measures precluded quantitative synthesis and meta-analysis, making a narrative approach more appropriate for integrating the existing evidence [18, 19].

2.2. Literature Search Strategy

A comprehensive literature search was conducted using five electronic databases: PubMed, Scopus, Web of Science, ScienceDirect, and Google Scholar. The search was performed to identify studies published between January 2014 and June 2026. Additional relevant publications published before 2014 were also considered when they provided foundational evidence regarding pharmacological mechanisms, phytochemical composition, or antihypertensive activity of *Anredera cordifolia*. The search strategy combined Medical Subject Headings (MeSH) and free-text terms using Boolean operators as follows:

("Anredera cordifolia" OR "Binahong") AND ("Hypertension" OR "High Blood Pressure") AND ("Herbal Medicine" OR "Medicinal Plant" OR "Complementary Therapy") AND ("Antihypertensive Effect" OR "Blood Pressure" OR "ACE Inhibition" OR "Vasodilation")

To ensure comprehensive coverage, supplementary searches were conducted using additional keywords related to phytochemical composition, antioxidant activity, endothelial function, vascular relaxation, and cardiovascular effects. Reference lists of eligible articles were manually screened to identify additional studies that may not have been retrieved through electronic database searches.

2.3. Eligibility Criteria

2.3.1. Inclusion Criteria

Studies were eligible for inclusion if they met the following criteria:

- Original research articles published in peer-reviewed journals.
- Investigations of *Anredera cordifolia* leaves, leaf extracts, leaf fractions, or leaf decoctions.
- Experimental laboratory studies evaluating antihypertensive mechanisms.
- In vitro studies assessing ACE inhibition, antioxidant activity, vasodilation, or endothelial effects.
- Ex vivo vascular studies.
- Animal studies reporting blood pressure outcomes.
- Human intervention studies involving participants with hypertension.
- Studies reporting cardiovascular, vascular, or blood pressure-related outcomes.

2.3.2. Exclusion Criteria

Studies were excluded if they met any of the following criteria:

- Editorials, commentaries, or opinion pieces.
- Conference abstracts without full-text availability.
- Duplicate publications.
- Studies unrelated to cardiovascular or antihypertensive effects.
- Articles lacking sufficient methodological detail.
- Studies focusing exclusively on non-leaf plant parts without cardiovascular relevance.

2.4. Study Selection Process

The titles and abstracts of identified records were screened for relevance. Potentially eligible articles underwent full-text assessment. Studies meeting the eligibility criteria were included in the narrative synthesis. The selection process prioritized studies directly evaluating antihypertensive activity. However, because direct clinical evidence

was limited, mechanistic studies investigating vasodilatory effects, ACE inhibition, antioxidant activity, endothelial protection, and anti-inflammatory properties were also included to provide a comprehensive understanding of the biological plausibility of antihypertensive effects.

For clarity of presentation, the included studies were grouped into two categories: (1) direct evidence evaluating *Anredera cordifolia* (Binahong), and (2) comparative evidence from other herbal antihypertensive plants that provided contextual support for the proposed biological mechanisms.

The study selection process followed the PRISMA 2020 recommendations. After removing duplicate records and screening titles and abstracts, potentially eligible articles underwent full-text review. The detailed process is presented in **Figure 1**. The study selection process followed a structured approach involving identification, screening, eligibility assessment, and final inclusion of relevant studies (**Figure 1**). Following the removal of duplicate records, all retrieved articles were screened based on their titles and abstracts to determine their relevance to the review objectives. Articles considered potentially eligible subsequently underwent full-text evaluation according to the predefined inclusion and exclusion criteria. To improve the transparency and reliability of the selection process, two independent reviewers independently screened all retrieved records by title, abstract, and full text. Any disagreements regarding study eligibility were resolved through discussion until consensus was achieved. Studies that failed to satisfy the eligibility criteria or did not provide sufficient information relevant to the review objectives were excluded from the final synthesis.




Figure 1. PRISMA 2020 flow diagram of literature selection process.

2.5. Scoring and Interpretation

Menstrual practice needs were assessed using the 36-item Menstrual Practice Needs Scale (MPNS-36) [17]. The MPNS-36 is designed to evaluate the extent to which menstrual management environments, resources, and experiences meet the needs of menstruating individuals across multiple domains, including access to menstrual materials, privacy, water and sanitation facilities, disposal options, comfort, and safety. Each item is rated on a four-point Likert scale reflecting the degree to which menstrual needs are met. Response options are scored from 0 to 3, with higher scores indicating that menstrual practice needs are better met. Negatively worded items are reverse-coded before analysis to ensure that all items are oriented in the same direction. After recoding, item scores are summed and averaged to generate an overall MPNS-36 score, with possible values ranging from 0 to

3. In the present study, the MPNS-36 score was analyzed as a continuous variable, with higher values representing better menstrual practice experiences and greater fulfillment of menstrual health and hygiene needs.

2.6. Data Extraction

Relevant information was extracted from each included study using a standardized evidence extraction framework. The following variables were collected:

- Author(s) and publication year
- Country of study
- Study design
- Experimental model or participant characteristics
- Type of Binahong preparation
- Intervention protocol
- Outcome measures
- Main findings
- Proposed antihypertensive mechanisms
- Study limitations

A total of 11 studies were identified as relevant to the antihypertensive potential of Binahong and were included in the evidence matrix. These studies comprised human intervention studies, animal experiments, ex vivo vascular investigations, phytochemical analyses, antioxidant studies, and pharmacological mechanism studies.

2.7. Data Synthesis

The extracted evidence was synthesized narratively and organized into thematic categories:

- Botanical characteristics of *Anredera cordifolia*.
- Phytochemical constituents.
- Mechanisms of antihypertensive action.
- Experimental evidence.
- Human evidence.
- Preparation and administration of Binahong leaf decoction.
- Safety and adverse effects.

To strengthen interpretation, studies were further categorized according to their contribution to the evidence base, namely direct antihypertensive evidence, mechanistic evidence, phytochemical evidence, and supportive cardiovascular evidence.

2.8. Quality Considerations

Given the narrative nature of this review, a formal risk-of-bias assessment was not conducted. Nevertheless, methodological quality was critically considered during evidence synthesis. Particular attention was paid to study design, sample size, intervention standardization, outcome measurement, duration of follow-up, and applicability to clinical hypertension management. The strength of evidence was interpreted cautiously because the majority of available studies were experimental or mechanistic in nature, whereas high-quality randomized controlled trials in humans remain scarce. Consequently, conclusions were drawn based on the overall consistency, biological plausibility, and methodological robustness of the available literature rather than on any single study.

2.9. Methodological Considerations

Although a formal risk-of-bias assessment was not performed because this study was designed as a narrative review, methodological quality was considered during data synthesis. Particular attention was given to study design, sample size, intervention characteristics, outcome measures, and the overall strength of the available evidence when interpreting the findings.

3. RESULTS

3.1. Literature Search Results

The literature search identified studies examining the antihypertensive potential of *Anredera cordifolia* (Binahong) through multiple research approaches, including phytochemical investigations, mechanistic pharmacology studies, ex vivo vascular experiments, animal studies, and human intervention studies. Following screening and eligibility assessment, 11 studies were considered relevant for inclusion in this narrative review.

The included studies represented a diverse body of evidence. Most studies focused on phytochemical characterization and mechanistic pathways, whereas only a limited number directly evaluated blood pressure outcomes in animal or human subjects. Consequently, the evidence base was categorized into four major domains: (i) direct antihypertensive evidence, (ii) mechanistic evidence, (iii) phytochemical evidence, and (iv) supportive cardiovascular evidence.

Only a limited number of studies have directly investigated the antihypertensive potential of *Anredera cordifolia* (**Table 1**). Most available evidence consists of phytochemical investigations, antioxidant studies, and mechanistic experiments demonstrating vasodilatory activity, moderate ACE inhibition, and endothelial-protective effects. These findings provide biological plausibility for the potential role of Binahong in blood pressure regulation but do not constitute conclusive clinical evidence.

Table 1. Summary of evidence on the antihypertensive potential of *Anredera cordifolia* (Binahong) and comparative herbal plants.

No.	Author (s), Years	Plant / Focus	Study Type	Sample / Model	Intervention / Assessment	Principal Findings
Direct evidence on <i>Anredera cordifolia</i> (Binahong)						
1	Astuti et al. 2011 [9]	<i>A. cordifolia</i> (leaves, stems, tubers)	Phytochemical study	Plant parts	Saponin analysis	In all plant parts have potential pharmacological activity relevant to cardiovascular health.
2	Djamil et al. 2012 [13]	Binahong leaf extract	Phytochemical and antioxidant study	Leaf extract	Flavonoid isolation and antioxidant assay	Flavonoids isolated with significant antioxidant activity.
3	Selawa et al. 2013 [12]	Ethanol extract of Binahong leaves	Phytochemical and antioxidant study	Leaf extract	Flavonoid and antioxidant analysis	High flavonoid content and strong antioxidant activity observed.
4	Garmana et al. 2017 [10]	Binahong leaf extract fractions	Experimental pharmacological study	Vascular tissue	Vasodilation and ACE inhibition assays	Demonstrated vasodilatory activity and moderate ACE inhibition supporting antihypertensive potential.
5	Susanti et al. 2024 [20]	Binahong extract	Phytochemical and antioxidant study	Plant extract	Phenolic and antioxidant evaluation	High phenolic content and antioxidant properties identified.
6	Sidhartha et al. 2024 [24]	Binahong leaf extract	Phytochemical, antioxidant and antifungal study	Leaf extract	Phytochemical screening and DPPH assay	Flavonoids, alkaloids, tannins and saponins with significant antioxidant activity.
7	Herawati et al. 2025 [14]	Ethanol extract of Binahong leaves	Phytochemical and antioxidant study	Leaf extract	Phenolic, flavonoid and antioxidant analysis	High phenolic and flavonoid levels associated with strong antioxidant potential.
Comparative evidence from other herbal antihypertensive plants						
8	Khayyal et al. 2019 [21]	<i>Olea europaea</i> (olive leaf)	Animal study	Hypertensive rats	Olive leaf extract	Reduced blood pressure and improved blood vessel function through antioxidant and endothelial protective effects, similar to Binahong.
9	Lockyer et al. 2011 [22]	<i>Olea europaea</i> (olive leaf)	Randomized controlled trial	Human participants	Olive leaf extract	Improved blood pressure and cardiovascular risk; supports clinical potential of phytochemical- rich herbal interventions.
10	Aekthammarat et al. 2024 [23]	<i>Moringa oleifera</i>	Animal study	L-NAME hypertensive rats	Leaf extract	Reduced blood pressure via antioxidant and endothelial protective mechanisms, similar to Binahong.
11	Susalit et al. [25]	<i>Olea europaea</i> (olive leaf)	Randomized clinical trial	Patients with stage-1 hypertension	Olive leaf extract vs. captopril	Antihypertensive efficacy comparable to captopril; provides comparative clinical evidence for herbal antihypertensive therapy.

To provide broader scientific context, comparative evidence from other phytochemical-rich medicinal plants was also reviewed (**Table 1**). Studies involving olive leaf (*Olea europaea*) and *Moringa oleifera* demonstrated significant antihypertensive effects through antioxidant activity, improved endothelial function, and modulation of vascular homeostasis. These studies were included solely as indirect evidence supporting the plausibility of

plant-derived antihypertensive mechanisms and should not be interpreted as direct evidence of the clinical efficacy of Binahong.

Table 2. Major phytochemical constituents of Binahong leaves and their potential relevance to hypertension.

Compound Group	Reported Biological Activity	Potential Relevance to Hypertension
Flavonoids	Antioxidant, endothelial protection	Enhancement of nitric oxide bioavailability and reduction of oxidative stress
Polyphenols	Antioxidant	Protection against vascular oxidative damage
Saponins	Anti-inflammatory, vasoprotective	Improvement of vascular function
Alkaloids	Cardiovascular modulation	Regulation of vascular tone
Terpenoids	Vasorelaxant activity	Support of vascular relaxation
Tannins	Antioxidant	Preservation of endothelial integrity
Glycosides	Cardiovascular activity	Potential modulation of vascular responses

As shown in **Table 2**, among these compounds, flavonoids and polyphenols were the most consistently reported constituents across studies. Quantitative analyses demonstrated substantial concentrations of these compounds in ethanol extracts of Binahong leaves. Antioxidant investigations conducted by Selawa et al. [12], Sidhartha et al. [20], and Herawati et al. [14] further demonstrated significant free-radical scavenging activity, supporting the potential role of Binahong in mitigating oxidative stress-related vascular dysfunction.

3.2. Mechanisms Potentially Relevant to Blood Pressure Regulation

The available mechanistic and phytochemical studies suggest several biological pathways through which the plant may influence cardiovascular function (**Table 3**). Experimental pharmacological studies reported that Binahong extracts exhibited vasodilatory activity in isolated vascular tissues and demonstrated moderate ACE inhibitory effects. These findings suggest that Binahong may influence blood pressure regulation through both vascular and neurohormonal pathways. In addition, the strong antioxidant properties consistently reported across phytochemical studies provide a biologically plausible mechanism by which Binahong may support endothelial function and vascular homeostasis.

Table 3. Proposed mechanisms relevant to the potential antihypertensive effects of Binahong.

Mechanism	Supporting Evidence	Potential Impact on Cardiovascular Function
Vasodilation	Garmana et al. [10]	Reduction of peripheral vascular resistance
ACE inhibition	Garmana et al. [10]	Potential attenuation of the renin-angiotensin system
Antioxidant activity	Djamil et al. [13]; Selawa et al. [12]; Sidhartha et al. [20]; Herawati et al. [14]	Reduction of oxidative stress and vascular damage
Endothelial protection	Flavonoid- and polyphenol-rich extracts	Preservation of endothelial function
Anti-inflammatory effects	Saponin- and polyphenol-containing extracts	Reduction of vascular inflammation

3.3. Evidence from Experimental and Comparative Studies

Only limited studies have directly investigated the antihypertensive effects of Binahong. Nevertheless, mechanistic evidence suggests that the plant possesses biological activities relevant to blood pressure regulation. Experimental work by Garmana et al. demonstrated vasodilatory effects and moderate ACE inhibitory activity of Binahong extracts, indicating potential cardiovascular benefits [10].

To provide contextual support for the biological plausibility of plant-based antihypertensive therapies, comparative evidence from other medicinal plants was also examined. Khayyal et al. (2002) reported significant blood pressure reduction and improved endothelial function following administration of olive leaf extract in hypertensive rats. Similarly, Aekthammarat et al. [21] demonstrated that *Moringa oleifera* leaf extract reduced blood pressure and alleviated vascular dysfunction in L-NAME-induced hypertensive rats through antioxidant and endothelial protective mechanisms. Human clinical evidence from Susalit et al. [22] further showed that olive leaf extract exhibited antihypertensive efficacy comparable to captopril among patients with stage-1 hypertension. Collectively, these findings suggest that phytochemical-rich medicinal plants may exert antihypertensive effects through multiple mechanisms involving antioxidant activity, vascular protection, and modulation of endothelial function. The presence of similar bioactive compounds in Binahong supports its potential role as a complementary intervention, although direct clinical evidence remains insufficient.

3.4. Preparation and Administration of Binahong Leaf Decoction

Traditional use of Binahong primarily involves the preparation of fresh leaves as a decoction (**Table 4**). Common preparation methods include washing mature leaves, boiling them in water for approximately 10–20 minutes, and consuming the resulting decoction after cooling. However, the reviewed studies revealed considerable variability

in extraction methods, solvent use, dosage, and duration of administration. The lack of standardized preparation protocols remains an important limitation and hinders direct comparison across studies.

Table 4. Characteristics of *Anredera cordifolia* (Binahong) preparations reported in the included studies.

Parameter	Observed Variation
Plant material	Fresh leaves, dried leaves, ethanol extracts
Preparation method	Decoction, extract, fraction
Administration route	Oral administration
Dosage	Not standardized
Treatment duration	Variable across studies

3.5. Safety and Potential Adverse Effects

According to **Table 5** safety data specific to Binahong remain limited. The available studies generally reported favorable tolerability and did not identify serious adverse effects associated with commonly used preparations. However, most investigations focused on phytochemical characterization and biological activity rather than comprehensive toxicological evaluation. Overall, the current literature suggests a favorable preliminary safety profile; however, rigorous toxicological and clinical safety studies remain necessary.

Table 5. Summary of available safety data for *Anredera cordifolia* (Binahong) preparations.

Safety Aspect	Current Evidence
Acute toxicity	Limited evidence suggests low toxicity
Serious adverse effects	Not reported in reviewed studies
Long-term safety	Insufficient evidence
Herb–drug interactions	Insufficient evidence
Use during pregnancy and lactation	Insufficient evidence

3.6. Overall Synthesis of Findings

As depicted in **Figure 2**, *Anredera cordifolia* (Binahong) contains several bioactive compounds, including flavonoids, polyphenols, saponins, alkaloids, terpenoids, and tannins, that exhibit antioxidant and vasoprotective properties. Current evidence suggests that these compounds may help regulate blood pressure by reducing oxidative stress, improving endothelial function, promoting vasodilation, and moderately inhibiting ACE. However, the evidence is strongest from phytochemical, mechanistic, and animal studies, whereas human

evidence remains limited and methodologically weak. Therefore, although Binahong shows promising potential as a complementary approach to hypertension management, well-designed preclinical studies and large randomized controlled trials are needed to confirm its efficacy and safety.




Figure 2. Proposed mechanisms and current evidence supporting the antihypertensive potential of *Anredera cordifolia* (Binahong).

4. DISCUSSION

This narrative review synthesized the available evidence regarding the potential role of *Anredera cordifolia* (Binahong) in hypertension management. The current literature suggests that Binahong possesses several biological properties that may contribute to blood pressure regulation, including antioxidant activity, vasodilatory effects ACE inhibition, endothelial protection, and anti-inflammatory activity [9-12, 20-24]. These findings provide biological credibility for the traditional use of Binahong as a complementary herbal therapy. However, the available evidence is not enough to proof of clinical antihypertensive efficacy because most studies have been conducted in laboratory settings or experimental models rather than in well-designed clinical trials [25, 26].

Hypertension is a multifactorial disorder driven by oxidative stress, endothelial dysfunction, chronic vascular inflammation, arterial remodeling, sympathetic overactivity, and dysregulation of the renin–angiotensin–aldosterone system [25-29]. These processes reduce nitric oxide bioavailability, impair vascular relaxation, and sustain elevated blood pressure. Binahong's phytochemical profile aligns closely with these pathways. Flavonoids and polyphenols can scavenge reactive oxygen species, preserve endothelial nitric oxide synthase activity, and enhance nitric oxide signaling [11-14, 29, 30]. Experimental work has further demonstrated direct vascular smooth-muscle relaxation and moderate ACE inhibition, suggesting complementary effects on both vascular tone and neurohormonal regulation [10]. Anti-inflammatory actions attributable to saponins and phenolic compounds may additionally attenuate vascular inflammation [9].

These mechanistic findings parallel those reported for better-characterized herbal antihypertensives. Olive leaf (*Olea europaea*) extract has been shown to lower blood pressure and improve endothelial function in both animal models and patients with stage-1 hypertension, with efficacy comparable to captopril in one randomized trial [21, 22]. *Moringa oleifera* leaf extract similarly reduces blood pressure in L-NAME-induced hypertensive rats through antioxidant and endothelial-protective mechanisms [21]. While such comparative data illustrate the therapeutic potential of phytochemical-rich plants, they cannot be extrapolated as direct evidence of Binahong's clinical effectiveness [23, 24, 31].

The current evidence base is constrained by several important limitations. High-quality human data are scarce. Most available investigations are phytochemical, in-vitro, ex-vivo, or animal studies [9-12, 20, 32]. The few human studies that have examined blood-pressure outcomes are small, quasi-experimental or pretest posttest designs that lack randomization, allocation concealment, blinding, and adequate control groups [15, 17, 33]. Short intervention periods further limit statistical power and increase susceptibility to selection bias, placebo effects, and regression to the mean. Marked heterogeneity in plant material (fresh versus dried leaves), extraction solvents, dosage regimens, treatment duration, and outcome measures precludes reliable cross-study comparison and

identification of optimal protocols [10, 12, 14]. Methodological reporting is frequently incomplete, leaving overall risk of bias uncertain. Publication bias may also inflate apparent benefits, as positive findings are more likely to appear in the literature. Long-term safety data including chronic toxicity, hepatotoxicity, nephrotoxicity, reproductive safety, and herb–drug interactions remain largely absent [16, 34, 35].

From a clinical perspective, Binahong cannot currently be recommended as an evidence-based antihypertensive therapy or as a substitute for conventional pharmacological treatment, lifestyle modification, or routine blood-pressure monitoring [23, 26]. It may be considered, under professional supervision, as a culturally acceptable complementary option for selected patients who wish to incorporate traditional medicine. Theoretical additive effects with ACE inhibitors, angiotensin-receptor blockers, calcium-channel blockers, beta-blockers, or diuretics warrant caution; patients should disclose herbal use and undergo regular monitoring until dedicated interaction studies become available [24, 35].

In resource-limited settings, Binahong's local availability and cultural acceptance may support its exploration as an adjunctive intervention. However, integration into public-health programs requires standardized preparations, quality control, pharmacovigilance, and robust clinical evidence rather than reliance on traditional use alone [16, 23].

Future research should prioritize multicenter randomized controlled trials that employ standardized Binahong preparations, adequate sample sizes, longer follow-up, blinded outcome assessment, and comprehensive evaluation of efficacy, safety, and potential herb-drug interactions. Parallel work on phytochemical standardization, dose response relationships, and long-term toxicology will be essential to determine whether Binahong can occupy a defined role within integrative hypertension care [26, 29, 36].

Strengths of the Current Evidence

The principal strength of the current evidence lies in the consistency of findings across phytochemical analyses, mechanistic pharmacology, ex vivo vascular experiments, and animal studies, all of which support the biological plausibility of Binahong as a complementary approach to hypertension management. The convergence of these independent lines of evidence provides a sound scientific rationale for future clinical investigation. Furthermore, the longstanding traditional use of Binahong offers valuable ethnopharmacological support for hypothesis generation, although traditional use alone cannot be considered evidence of clinical effectiveness.

Weaknesses and Limitations of Existing Studies

Till now several critical limitations substantially weaken confidence in the current evidence base. High-quality human data are scarce; most findings derive from laboratory, phytochemical, and animal studies rather than rigorously designed randomized controlled trials in hypertensive patients. The few available human studies are typically small, quasi-

experimental, and lack randomization, allocation concealment, blinding, or adequate control groups, thereby increasing susceptibility to selection, performance, and measurement bias. Short intervention periods further limit statistical power and generalizability. Marked heterogeneity in plant material, extraction methods, dosages, treatment duration, and outcome measures precludes meaningful comparison across studies and prevents identification of optimal therapeutic protocols. Methodological reporting is frequently incomplete, leaving overall risk of bias uncertain, while the possibility of publication bias may inflate apparent benefits. Finally, long-term safety data including chronic toxicity, organ-specific effects, reproductive safety, and herb-drug interactions remain largely absent.

Future Research Directions

Future investigations should prioritize multicenter randomized controlled trials with adequate sample sizes, concealed allocation, blinded outcome assessment, standardized Binahong preparations, and longer follow-up periods. Such studies should evaluate clinically relevant outcomes including systolic and diastolic blood pressure, endothelial function, cardiovascular biomarkers, quality of life, adverse events, and potential herb drug interactions.

Future systematic reviews should also assess the methodological quality of available studies using standardized risk-of-bias assessment tools and evaluate the potential influence of publication bias. Standardization of Binahong preparation methods, phytochemical characterization, dosage regimens, and quality control procedures will be essential for generating reproducible, clinically meaningful, and internationally comparable evidence. Furthermore, long-term studies evaluating chronic safety, hepatotoxicity, nephrotoxicity, and pharmacokinetic interactions with conventional antihypertensive medications are needed before definitive clinical recommendations can be established.

5. CONCLUSIONS

This narrative review indicates that *Anredera cordifolia* (Binahong) possesses phytochemical constituents and pharmacological properties that provide a biologically plausible basis for supporting blood pressure regulation. Current evidence demonstrates antioxidant, vasodilatory, endothelial-protective, anti-inflammatory, and moderate ACE inhibitory activities. However, the available evidence is derived predominantly from phytochemical investigations, mechanistic studies, and animal experiments, whereas robust clinical evidence in humans remains limited. Accordingly, Binahong cannot yet be recommended as an evidence-based antihypertensive therapy or as a substitute for standard pharmacological treatment. Nevertheless, it may represent a promising complementary intervention that warrants further investigation. Future multicenter randomized controlled trials using standardized Binahong preparations, adequate sample sizes, longer follow-up periods, and

comprehensive safety assessments are essential to establish its clinical efficacy, optimal dosage, and role in integrative hypertension care.

ACKNOWLEDGEMENT

The authors sincerely express their gratitude to their colleagues and affiliated institutions for their continuous support and encouragement throughout the research process.

FUNDING STATEMENT

This research received no specific grant from any funding agency, commercial, or not-for-profit sectors.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest related to this manuscript.

ETHICAL CONSIDERATION

Ethical approval was not required for this study because it is a narrative review based exclusively on previously published literature. No human participants, animals, or identifiable personal data were involved, and no primary data were collected.

AUTHOR CONTRIBUTIONS

La Ode Alifariki: Conceptualization, methodology, investigation, data curation, formal analysis, writing original draft, review, and supervision. **Dewi Nopiska Lilis:** methodology, data curation, review manuscript. all authors have read and approved the final manuscript and agree to be accountable for all aspects of the work

DECLARATION OF AI USE

During the preparation of this manuscript, the authors used artificial intelligence (AI)-assisted language tools solely to improve grammar, language clarity, and readability. The AI tools were not used to generate scientific content, interpret data, draw conclusions, or replace the authors' intellectual contributions. All literature selection, critical appraisal, interpretation of the evidence, and final manuscript preparation were performed entirely by the authors, who accept full responsibility for the accuracy and integrity of the manuscript.

REFERENCES

1. Mills KT, Stefanescu A, He J. The global epidemiology of hypertension. *Nat Rev Nephrol.* 2020;16(4):223–37. doi:10.1038/s41581-019-0244-2
2. Zhou B, Carrillo-Larco RM, Danaei G, Riley LM, Paciorek CJ, Stevens GA, et al. Worldwide trends in hypertension prevalence and progress in treatment and control from 1990 to 2019: a pooled analysis of 1201 population-representative studies with 104 million participants. *Lancet.* 2021;398(10304):957–80.
3. World Health Organization. Global Report on Hypertension 2025: High Stakes Turning Evidence Into Action. World Health Organization; 2025. Last visited 20 June, 2026.
4. Carey RM, Muntner P, Bosworth HB, Whelton PK. Reprint of: prevention and control of hypertension: JACC Health Promotion Series. *J Am Coll Cardiol.* 2018;72(23 Part B):2996–3011.
5. Carey RM, Whelton PK. 2017 ACC/AHA Hypertension Guideline Writing Committee. Prevention, detection, evaluation, and management of high blood pressure in adults: synopsis of the 2017 American College of Cardiology/American Heart Association Hypertension Guideline. *Ann Intern Med.* 2018;168(5):351–358
6. Montezano AC, Touyz RM. Reactive oxygen species and endothelial function—role of nitric oxide synthase uncoupling and Nox family nicotinamide adenine dinucleotide phosphate oxidases. *Basic Clin Pharmacol Toxicol.* 2012;110(1):87–94.
7. Abplanalp WT, John D, Cremer S, Assmus B, Dorsheimer L, Hoffmann J, et al. Single-cell RNA-sequencing reveals profound changes in circulating immune cells in patients with heart failure. *Cardiovasc Res.* 2021;117(2):484–94.
8. Pertiwi IA. *Pengaruh Terapi Komplementer Air Rebusan Daun Binahong terhadap Tekanan Darah pada Lansia dengan Hipertensi di RW 03 Kelurahan Limo Kota Depok.* Bachelor's thesis. Universitas Pembangunan Nasional Veteran Jakarta; 2017.
9. Astuti SM, Sakinah MAM, Andayani RBM, Risch A. Determination of saponin compound from *Anredera cordifolia* (Ten.) Steenis plant (Binahong) to potential treatment for several diseases. *J Agric Sci.* 2011;3(4):224.
10. Garmana AN, Sukandar EY, Fidrianny I. Efek Vasodilatasi dan Inhibisi Angiotensin Converting Enzyme dari Ekstrak Etanol dan Fraksi Daun Binahong (*Anredera Cordifolia* (Ten). V. Steenis). *Acta Pharm Indones [Internet].* 2017;42(2):51–9.
11. Fraga CG, Croft KD, Kennedy DO, Tomás-Barberán FA. The effects of polyphenols and other bioactives on human health. *Food Funct.* 2019;10(2):514–28.
12. Rusdiana, Widyawati T, Sari DK, Widjaja SS. Phytochemical analysis of the ethanol extract of *Anredera cordifolia* (Ten.) Steenis leaves by UV-Vis spectroscopy. *Baghdad Sci J.* 2024;21(11):3446–3451. doi:10.21123/bsj.2024.9354.
13. Djamil R, Wahyudi PS, Wahono S, Hanafi M. Antioxidant activity of flavonoid from *Anredera cordifolia* (Ten) Steenis leaves. *Int Res J Pharm.* 2012;3(9):241–3.
14. Herawati IE, Inayah I, Dewi L. Kadar Fenolik Total, Flavonoid Total, Dan Aktivitas Antioksidan Ekstrak Etanol Daun Binahong (*Anredera cordifolia* Ten. Steenis). *JIFI (Jurnal Ilm Farm Imelda).* 2025;9(1):8–16.
15. Wijayanti ES, Kurniawan WE. Pengaruh Pemberian Air Rebusan Daun Binahong (*Anredera Cordifolia*) Terhadap Tekanan Darah Pada Penderita Hipertensi di Wilayah Desa Jetis Kecamatan Nusawungu. *SENTRI J Ris Ilm.* 2023;2(3):671–81.
16. Ekor M. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. *Front Pharmacol.* 2014;5:177.
17. Firdaus M, Rezeki NT. Pengaruh Mengkonsumsi Air Rebusan Daun Binahong (*Anredera Cordifolia*) terhadap Penurunan Tekanan Darah Tinggi pada Lansia di UPT PSTW Khusnul Khotimah Pekanbaru. *J STIKes Awal Bros pekanbaru.* 2020;1(2):29–37.
18. Green BN, Johnson CD, Adams A. Writing narrative literature reviews for peer-reviewed journals: secrets of the trade. *J Chiropr Med.* 2006;5(3):101–17.
19. Ferrari R. Writing narrative style literature reviews. *Med Writ.* 2015;24(4):230–5.
20. Sidhartha E, Yang JJ, Dimara RSN, Rahmawati F, Purba SWD. Phytochemical screening, antioxidant and antifungal activity test of Binahong leaf extract (*Anredera cordifolia* (Ten.) Steenis). *Eur J Adv Chem Res.* 2024;5(1):1–8.
21. Aekthammarat D, Pannangpetch P, Tangsucharit P. Moringa oleifera leaf extract lowers high blood pressure by alleviating vascular dysfunction and decreasing oxidative stress in L-NAME hypertensive rats. *Phytomedicine.* 2019;54:9–16.
22. Susalit E, Agus N, Effendi I, Tjandrawinata RR, Nofiarny D, Perrinjaquet-Moccetti T, et al. Olive (*Olea europaea*)

- leaf extract effective in patients with stage-1 hypertension: comparison with Captopril. *Phytomedicine*. 2011;18(4):251–8.
23. Choi D, Im H Bin, Choi SJ, Han D. Safety classification of herbal medicine use among hypertensive patients: a systematic review and meta-analysis. *Front Pharmacol*. 2024;15:1321523. doi:10.3389/fphar.2024.1321523
 24. Nyulas KI, Simon-Szabó Z, Pál S, Fodor MA, Dénes L, Cseh MJ, et al. Cardiovascular Effects of Herbal Products and Their Interaction with Antihypertensive Drugs-Comprehensive Review. *Int J Mol Sci*. 2024;25(12). doi:10.3390/ijms25126388
 25. Rodrigo R, González J, Paoletto F. The role of oxidative stress in the pathophysiology of hypertension. *Hypertens Res*. 2011;34(4):431–40. doi:10.1038/hr.2010.264
 26. Chowdhury S, Mitra P, Chakraborty P, Kar K. Herbal antihypertensives: Integrative perspectives from traditional medicine and clinical pharmacology up to 2025. *Clin Tradit Med Pharmacol*. 2025;6(4):200246. doi:https://doi.org/10.1016/j.ctmp.2025.200246
 27. Madamanchi NR, Runge MS. Redox signaling in cardiovascular health and disease. *Free Radic Biol Med*. 2013;61:473–501.
 28. Jeremic J, Govoruskina N, Bradic J, Milosavljevic I, Srejavic I, Zivkovic V. Sacubitril/valsartan reverses cardiac structure and function in experimental model of hypertension-induced hypertrophic cardiomyopathy. *Mol Cell Biochem*. 2023;478(12):2645–2656. doi: 10.1007/s11010-023-04690-7.
 29. Harrison DG, Guzik TJ, Lob HE, Madhur MS, Marvar PJ, Thabet SR, et al. Inflammation, immunity, and hypertension. *Hypertension*. 2011;57(2):132–40.
 30. Fitria VA, Sakdiyah AK, Kurniati V, Putra DOY, Putri NZ, Pratiwi PD. Profil Farmakologi Binahong (*Anredera cordifolia*) Sebagai Kandidat Modern Medicine: Tinjauan Sistematis. *Sinteza*. 2025; 28;5:32–47. doi:10.29408/sinteza.v5i1.29368
 31. Susanti H. Total phenolic content and antioxidant activities of Binahong (*Anredera cordifolia*.). *JKKI J Kedokt dan Kesehat Indones*. 2019;171–5.
 32. Mandol A, Anjum S, Kayes MN, Khan SN, Khatun Z, Parvin R, Haque S. Sun protection factor (spf) assessment of commercial sunscreens in vitro and sunscreen awareness among university students in bangladesh. *J. Biosci. Public Health*. 2026;2(3):264–277. doi:10.5455/JBPH.2026.12
 33. Jin Z, Lan Y, Li J, Wang P, Xiong X. The role of Chinese herbal medicine in the regulation of oxidative stress in treating hypertension: from therapeutics to mechanisms. *Chin Med*. 2024;19(1):150.
 34. Maelani I, Srihandayani N, Maulana F, Syahbani MW, Ramdhani R, Mishel S, et al. Pengaruh Daun Binahong Terhadap Penurunan Tekanan Darah. *J Nurs Pract Sci*. 2023;2(1):172–7.
 35. Vanhoutte PM, Shimokawa H, Feletou M, Tang EHC. Endothelial dysfunction and vascular disease: a 30th anniversary update. *Acta Physiol*. 2017;219(1):22–96.
 36. Shrivastav G, Dubey A, Kumari M, Shukla T. Advances in pharmacognosy for modern drug discovery and development. *J. Biosci. Public Health*. 2025;1(3):45–63. doi:10.5455/JBPH.2025.14